

On the Absence of the Zeno Effect in Relativistic Quantum Field Theory

Ramón F. Alvarez-Estrada*

Departamento de Física Teórica, Universidad Complutense, 28040 Madrid, Spain

José L. Sánchez-Gómez†

Departamento de Física Teórica, Universidad Autónoma de Madrid, 28049-Madrid, Spain

February 1, 2008

Abstract

We study the time evolution of decaying particles in renormalizable models of Relativistic Quantum Field Theory. Significant differences between the latter and Non Relativistic Quantum Mechanics are found—in particular, the Zeno effect seems to be absent in such RQFT models. Conventional renormalization yields finite time behaviour in some cases but fails to produce finite survival probabilities in others.

The analysis of the finite time evolution in relativistic quantum field theory (RQFT) and, in particular, the behaviour at short times of the survival probability of unstable particles in RQFT is a relevant matter as far as the foundations of quantum mechanics are concerned and has some peculiarities which make it to be worth studying. But before going to RQFT, we will briefly recapitulate the main features of the survival amplitude (SA) at short times in non relativistic quantum mechanics (QM henceforth); see, for instance, [1] for a comprehensive review.

*E-mail: ralvarez@eucmax.sim.ucm.es

†E-mail: jolu@delta.ft.uam.es

Let $P(t)$ ($t > 0$) the survival probability (SP) for an unstable system represented in QM by some normalized state $|\Psi\rangle$, and let H be the Hamiltonian governing the evolution of the quantum system. The SP is given by $P(t) = |\langle \Psi | \exp(-itH) | \Psi \rangle|^2$ ($\hbar = 1$). Then, assuming the finiteness of the energy dispersion in the initial (unstable) state, i.e.,

$$(\Delta E)^2 = \langle \Psi | H^2 | \Psi \rangle - (\langle \Psi | H | \Psi \rangle)^2 < \infty, \quad (1)$$

as indeed happens in QM, one can show that at very short times

$$P(t) \simeq 1 - (\Delta E)^2 t^2 \quad (t \rightarrow 0^+). \quad (2)$$

Notice that the last equation shows the violation of the exponential decay law at very short times. The quadratic behaviour in t gives rise to the so-called quantum Zeno effect, whose consequences and possible experimental detection [2] have been profusely discussed (see [3] and references therein for some recent work on this effect).

Another interesting point — which, however, is not the main subject of this paper — is the behaviour of the survival probability for large t values. One knows that if the hamiltonian spectrum is bounded from below, then the decay cannot be exponential, and $P(t) > P_{exp}(t)$, as $t \rightarrow \infty$. Now, whereas that property (positivity of the energy) always holds in QM, the situation is less clear in RQFT. Let H_{ren} be the hamiltonian of a certain model of RQFT, after having carried out standard ultraviolet renormalization. To the best of our knowledge, H_{ren} has been *rigorously* constructed for superrenormalizable models, like, for instance, the $\lambda\phi^4$ and Yukawa RQFT, in 1+1 dimensions, but presumably not for renormalizable models of the kind dealt with in this work. In the superrenormalizable RQFT models for which H_{ren} exists, the latter is lower bounded [4]. In any case, it is not the lower boundedness of the hamiltonian but, rather, the finiteness of the energy dispersion in the initial (unstable) state what is relevant for the existence of the Zeno effect.

Regarding the RQFT case, it has been pointed out [5] that the unrenormalized survival amplitude of an unstable particle has, in any model of RQFT (superrenormalizable or renormalizable), a singular behaviour as $t \rightarrow 0^+$ and, in order to provide a cure for it, a new characteristic time has been introduced. Such singularities seem to be related to those already discussed

by Bogoliubov and Shirkov [6] for the Schrödinger state vector in RQFT. More recently, the present authors have shown that in a superrenormalizable model of RQFT (namely, a direct extension of the scalar ϕ^3 -theory to the decay of an unstable particle) the above short-time singularities of the survival amplitude can be fully eliminated through a conventional renormalization procedure [7]. Here, we are going to analyze the time evolution of the SP of unstable particles in renormalizable models of RQFT. Now, one should be careful about vacuum polarizing interactions and ultraviolet divergent renormalization. Let us clearly state the problem

We consider —with respect to a given reference frame— some “free” Hamiltonian H_0 which describes particles for $t \leq 0$ in RQFT in the Schrödinger picture. Also, let $\vec{P}$ the total three-momentum operator, with $[\vec{P}, H_0] = 0$, and let $|i, \vec{p}_i\rangle$ the initial state representing the unstable particle at $t = 0$, with four-momentum $p_i = (E_i, \vec{p}_i)$, $E_i = (m_i^2 + \vec{p}_i^2)^{1/2}$, and renormalized mass m_i . Thus, $|i, \vec{p}_i\rangle$ is a common eigenstate of both H_0 and $\vec{P}$. If $|0\rangle$ represents the vacuum of H_0 , then

$$|i, \vec{p}_i\rangle \equiv |i\rangle = a^+(\vec{p}_i) |0\rangle, \quad (3)$$

where $a^+(\vec{p}_i)$ is the associated creation operator (in the Schrödinger picture). The interaction Hamiltonian H_I , which acts for $t > 0$ and is responsible for the decay has, by assumption, the following properties:

1. It commutes with $\vec{P}$.
2. It gives rise to a renormalizable local RQFT.
3. It involves a small coupling constant, in the sense that one can reliably apply perturbation theory.

Upon going over to the interaction picture (ip) and after some straightforward calculations, we get the basic formula enabling us to compute the survival amplitude (SA) in renormalized perturbation theory

$$A(t) = \langle i | i \rangle^{-1} \sum_{n=0}^{\infty} \frac{(-i)^n}{n!} \int_0^t dt_1 \cdots \int_0^t dt_n \langle 0 | T[a(t)H_I(t_1) \dots H_I(t_n)a^+(0)] | 0 \rangle \quad (4)$$

where, for simplicity, we have omitted i in the operators, as well as $\vec{p}_i$ in a and a^+ . Of course, T stands for time-ordered product. One can now evaluate $A(t)$ by using (4) and Wick's theorem in the standard way. Then it will be useful to introduce the following factorized form of the SA

$$A(t) = \exp(-iE_i t) \tilde{A}(t), \quad (5)$$

where the “reduced” survival amplitude $\tilde{A}(t) \rightarrow 1$ for any $t > 0$ if $H_I \rightarrow 0$. Now we have to deal with the divergences generically known as disconnected vacuum contributions (dvc). For the sake of brevity, we do not present here the details of the calculation (see [7] for a more elaborated treatment), which yields the following result

$$\tilde{A}(t) = A_{dvc}(t) A_{ph}(t). \quad (6)$$

$A_{ph}(t)$, to be called hereafter the physical survival amplitude, is the sum of all perturbative contributions to $\tilde{A}(t)$ in each of which $a(t)$ is not contracted with $a^+(t=0)$. Therefore all the contributions to A_{ph} are free of infinite volume divergences; all the latter are included in the factor $A_{dvc}(t)$. We should notice that $A_{ph} \rightarrow 1$ as $H_I \rightarrow 0$. We shall concentrate on A_{ph} . As already said, the above procedure has been applied to the case of a superrenormalizable (ϕ^3 -like) interaction [7], having shown that, after a standard ultraviolet mass renormalization, the survival amplitude is finite for all t and it behaves linearly in t as $t \rightarrow 0$ (no Zeno effect, then). Nevertheless, the scope of such superrenormalizable interactions is rather limited, so that, here, we are going to deal with the case of renormalizable interactions, which is more relevant from a physical point of view.

We shall consider a (unstable) relativistic spin 1/2 particle (i), with renormalized mass m_i , which can decay into two relativistic particles (a , b), with renormalized masses m_a , m_b , so that: i) $m_i > m_a + m_b$, and ii) b is a spin-1/2 fermion while a is a scalar spinless boson (we are considering all the particles neutral for simplicity; the general case is physically equivalent). Now H_0 is the sum of three free Hamiltonians: one, corresponding to particle a being similar to those in the superrenormalizable case (kinetic energy of a scalar spin-0 boson) and the other two, of particles i and b , being the well-known for relativistic free Dirac fermions. For $t > 0$ the interaction Hamiltonian in

the Schrödinger picture is

$$H_I = \lambda \int d^3\vec{x} [\bar{\Psi}_b(x)\Psi_i(x)\Phi_a(x)] + \text{h.c.} - \delta m_i \int d^3\vec{x} N[\bar{\Psi}_i(x)\Psi_i(x)] - (Z_{i,2} - 1) \int d^3\vec{x} [\bar{\Psi}_i(\gamma \cdot \partial)\Psi_i - m_i \bar{\Psi}_i\Psi_i] + \text{o.c.t.} \quad (7)$$

where λ is a renormalized dimensionless coupling constant. The mass renormalization counterterm for the unstable i-fermion is, to order λ^2

$$\delta m_i = \frac{i\lambda^2}{(2\pi)^4} \int d^4k \frac{-k \cdot \gamma + m_b}{(k^2 - m_a^2 + i\epsilon)(k^2 - m_b^2 + i\epsilon)} \quad (8)$$

The term proportional to $Z_{i,2} - 1$ is the standard contribution corresponding to the wave-function renormalization for particle i , and o.c.t. denote other renormalization counterterms which are not relevant here. The calculation now turns out to be much more elaborated than in the superrenormalizable case. One has, to order λ^2 ,

$$A_{ph}(t) - 1 = \frac{m_i}{E_i} \bar{u}(\vec{p}_i, \sigma) K_{ren} u(\vec{p}_i, \sigma) \quad (9)$$

$$K_{ren} = K + i\delta m_i t + i(Z_{i,2} - 1)(\gamma \cdot p_i - m_i)t \quad (10)$$

$$K = \frac{\lambda^2}{(2\pi)^5} \int d^4k_1 d^4k_2 \frac{\gamma \cdot k_2 + m_b}{(k_1^2 - m_a^2 + i\epsilon)(k_2^2 - m_b^2 + i\epsilon)} \delta^{(3)}(\vec{k}_1 + \vec{k}_2 - \vec{p}_i) \cdot \frac{\exp[it(E_i - k_1^0 - k_2^0)] - 1}{i(E_i - k_1^0 - k_2^0)} \cdot \frac{\exp[it(k_1^0 + k_2^0 - E_i)] - 1}{i(k_1^0 + k_2^0 - E_i)}, \quad (11)$$

where $\epsilon \rightarrow 0^+$ and $u, \bar{u}$ denote normalized Dirac spinors for the unstable particle with polarization σ . In K , we integrate over $\vec{k}_2$ and, by adding $-i\epsilon$ to E_i (which is a consistent procedure), we perform a residue integration over k_2^0 . These integrations give K as a piece linear in t plus another contribution, Ξ . In Ξ , we integrate over k_1^0 by residues, then obtaining

$$K_{ren} = it\Sigma_{ren} + \lambda^2\Xi \quad (12)$$

$$\Sigma_{ren} - [\delta m_i + (\gamma \cdot p_i - m_i)(Z_{i,2} - 1)] = -i \frac{\lambda^2}{(2\pi)^4} \int \frac{d^4k_1}{k_1^2 - m_a^2 + i\epsilon} \cdot \frac{\gamma \cdot (p_i - k_1) + m_b}{(p_i - k_1)^2 - m_b^2 + i\epsilon} \equiv \Delta + iAbs\Sigma_{ren} \quad (13)$$

$$\Xi = \int \frac{d^3 \vec{k}_1}{(2\pi)^3} \frac{1}{4E_a E_b} \left\{ \frac{[\exp it(E_i - E_a - E_b) - 1][E_b \gamma^0 + (\vec{k}_1 - \vec{p}_i) \cdot \vec{\gamma} + m_b]}{(E_a + E_b - E_i - i\epsilon)^2} + \frac{[\exp it(E_i + E_a + E_b) - 1][-E_b \gamma^0 + (\vec{k}_1 - \vec{p}_i) \cdot \vec{\gamma} + m_b]}{(E_a + E_b + E_i - i\epsilon)^2} \right\} (14)$$

where $E_a = (m_a^2 + \vec{k}_1^2)^{1/2}$, $E_b = [m_b^2 + (\vec{p}_i - \vec{k}_1)^2]^{1/2}$.

Ξ is the sum of two contributions each of which seems logarithmically ultraviolet divergent through naive power counting. Fortunately, Ξ is finite for any $t \geq 0$, since each divergence proportional to $\vec{\gamma}$ cancels out by symmetric integration, while those proportional to γ^0 cancel out with one another. Σ_{ren} is the renormalized self-energy of the unstable particle. The quantities Δ and $Abs\Sigma_{ren}$ (the absorptive part of Σ_{ren}), defined uniquely through standard methods, are finite. We shall define the quantity Ξ_1 out of the first term on the right hand side of Eq. (14) as follows: i) we replace $\exp it(E_i - E_a - E_b) - 1$ by $i \sin t(E_i - E_a - E_b)$, ii) if $A \equiv E_a + E_b - E_i$, we replace $(A - i\epsilon)^{-2}$ by $-i\pi\delta'(A)$. (Recall that $(A - i\epsilon)^{-2} = PA^{-2} - i\pi\delta'(A)$, where P is the principal part and δ' stands for the derivative of the δ function.) This gives $\Xi = \Xi_1 + \Xi_2$, where Ξ_2 embodies all the remaining contributions to Ξ . Some direct computations yield the following properties:

1) After cancelling out a linearly ultraviolet divergent piece proportional to $\vec{\gamma}$ (by symmetric integration), Δ bears the form $\Delta_1 + \gamma^\mu \Delta_{2,\mu}$, Δ_1 and $\Delta_{2,\mu}$ being logarithmically divergent quantities.

2) By employing δm_i and the corresponding logarithmically ultraviolet divergent expression for $Z_{i,2} - 1$ to order λ^2 (see, for instance, [8]), all logarithmically ultraviolet divergences cancel out in $X \equiv \delta m_i + (\gamma \cdot p_i - m_i)(Z_{i,2} - 1) + \Delta$, which becomes finite. Moreover $\bar{u}(\vec{p}_i, \sigma) X u(\vec{p}_i, \sigma)$ is real.

3) Ξ_1 is linear and t , and, moreover, $-t Abs\Sigma_{ren} + \lambda^2 \Xi_1 = 0$

4) Ξ_2 is ultraviolet finite

We shall define Ξ_3 out of Ξ_2 as follows: in the expression yielding the latter, we replace $\exp it[E_i \mp (E_a + E_b)] - 1$ by $\cos t[E_i \mp (E_a + E_b)] - 1$. Then

we have $\Xi_2 = \Xi_3 + \Xi_4$, where

$$\begin{aligned} \Xi_3 = & \frac{1}{(2\pi)^3} \int \frac{d^3\vec{k}_1}{4E_a E_b} \left\{ \frac{\cos t(E_i - E_a - E_b) - 1}{(E_a + E_b - E_i)^2} [E_b \gamma^0 + (\vec{k}_1 - \vec{p}_i) \cdot \vec{\gamma} + m_b] + \right. \\ & \left. \frac{\cos t(E_i + E_a + E_b) - 1}{(E_a + E_b + E_i)^2} [-E_b \gamma^0 + (\vec{k}_1 - \vec{p}_i) \cdot \vec{\gamma} + m_b] \right\} \end{aligned} \quad (15)$$

while Ξ_4 contains all the remaining contributions to Ξ_2 . Then

$$K_{ren} = itX + \lambda^2 \Xi_4 + \lambda^2 \Xi_3$$

The key reasons for the successive definitions of Δ , Ξ_j , $j = 1, 2, 3, 4$ are the above results 1)–4) plus the following ones:

5) $\bar{u}(\vec{p}_i, \sigma)[itX + \lambda^2 \Xi_4]u(\vec{p}_i, \sigma)$ is purely imaginary,

6) $\bar{u}(\vec{p}_i, \sigma)\lambda^2 \Xi_3 u(\vec{p}_i, \sigma)$ is real

Consequently, the physical survival probability up to order λ^2 is

$$P(t) = |A_{ph}(t)|^2 = 1 + 2\lambda^2 [\bar{u}(p_i, \sigma) \Xi_3 u(\vec{p}_i, \sigma)] \frac{m_i}{E_i}. \quad (16)$$

We shall now deal with a very important physical requirement, namely, the consistency between (16) and Fermi's golden rule. For that purpose we let $t \rightarrow \infty$ and recall that, then, $q^{-2}[1 - \cos(tq)] \rightarrow \pi t \delta(q)$ for any real q . A glance at Eq. (15) shows that only the first fraction on its r.h.s. yields a non-vanishing contribution as $t \rightarrow \infty$. Now, using well-known properties of Dirac matrices and spinors, one readily gets

$$P(t) \simeq 1 - \frac{m_i}{E_i} \Gamma t \quad (t \rightarrow \infty), \quad (17)$$

where Γ can be written as

$$\Gamma = \frac{\lambda^2}{16\pi^2 m_i} \int \frac{d^3\vec{k}_1 d^3\vec{k}_2}{E_a(\vec{k}_1) E_b(\vec{k}_2)} (p_i k_2 + m_i m_b) \delta^4(p_i - k_1 - k_2). \quad (18)$$

Notice that Γ is relativistically invariant and coincides exactly (as a straightforward computation employing Dirac matrices and their traces show) with

the total decay rate of the unstable particle in its rest frame ($\vec{p}_i = 0$). By taking into account that $m_i/E_i = (1 - \vec{v}_i^2)^{1/2}$, $\vec{v}_i$ being the velocity of the decaying particle, and introducing its lifetime $\tau = \Gamma^{-1}$ at rest, Eq. (17) can also be cast as $P(t) = 1 - t(1 - \vec{v}_i^2)^{1/2}/\tau$, which is simply showing the relativistic fact that an unstable particle with velocity $\vec{v}_i$ in an inertial frame has a lifetime $\tau(1 - \vec{v}_i^2)^{-1/2}$ in that frame. Then, Eqs. (17-18) are consistent with the linearization of the exponential decay behaviour $\exp[-t(1 - \vec{v}_i^2)^{1/2}\Gamma]$ in an inertial frame; Γ , the decay rate in the rest frame, being given by Fermi's golden rule

Next, we shall investigate the behaviour of the survival probability in the rest frame ($\vec{p}_i = 0$) for small positive values of t . For that purpose, we write $\Xi_3 = Y_+ + \gamma^0 Y_-$ (noticing that the contribution proportional to $\vec{\gamma}$ vanishes by symmetric integration). We are interested in the behaviours of $Y_{\pm}$ for small positive t . Such behaviours are determined by large values of $|\vec{k}_1|$. It suffices to neglect all masses in Y_+ , and one readily obtains $Y_+ \sim -m_b t/(16\pi)$. The treatment of Y_- requires more care, due to an additional factor $E_b(\vec{k}_1)$ which appears with opposite signs in both contributions in the r.h.s. of (15) and, hence, leads to cancellations at large $|\vec{k}_1|$. One finds $Y_- \sim -tm_i/(32\pi)$. Then, for very small values of t , the survival probability is given by

$$P(t) \simeq 1 - \frac{\lambda^2}{16\pi} (2m_b + m_i)t \quad (19)$$

The linear behaviour for very small values of t appearing in (19) shows that the Zeno effect is not present.

We shall now consider another renormalizable model, similar regarding physical description and notations to the one previously described, except for the following crucial differences:

- a) The unstable particle i is a spinless boson which decays into two spin 1/2 (Dirac) fermions (a , b) (again, all particles are neutral for simplicity).
- b) For $t > 0$, the interaction hamiltonian is

$$H_I = \lambda \int d^3(\vec{x}) [\bar{\Psi}_a(\vec{x}) \Psi_b(\vec{x}) \Phi_i(\vec{x})] + \text{h.c.} + \text{c.t.} \quad (20)$$

The counterterms (c.t.) include both the mass and wave function renormalization counterterms for the decaying boson (see, for instance, [9] for their detailed expressions) plus another renormalization counterterms which are not relevant for our calculation. We shall limit ourselves to study the physical survival amplitude $A_{ph}(t)$ for the decaying boson in its rest frame ($\vec{p}_i = 0$). A computation entirely similar to the one carried out for the previous model yields

$$A_{ph}(t) \simeq 1 + \frac{1}{2m_i}(iXt + \lambda^2\Xi_4 + \lambda^2\Xi_3). \quad (21)$$

The quantities X and Ξ_4 are the respective analogues of those for the previous model: both iX and Ξ_4 are pure imaginary. X is the ultraviolet finite contribution which results after cancellations between: i) divergent terms arising from the actual analogue of K (Eq. (11) for the previous model) and the mass and wavefunction renormalization counterterms for the decaying boson, ii) finite contributions linear in t (analogues of $-t \text{ Abs}\Sigma_{ren}$ and $\lambda^2\Xi_1$ in the previous model). We shall concentrate in the relevant piece, which is real and, hence, contributes to the survival probability $P(t) = |A_{ph}(t)|^2$:

$$\begin{aligned} \Xi_3 = & \frac{1}{(2\pi)^3} \int \frac{d^3\vec{k}_1}{4E_a E_b} (m_a m_b + E_a E_b + \vec{k}_1^2) \cdot \\ & \left[\frac{\cos t(E_a + E_b - m_i) - 1}{(E_a + E_b - m_i)^2} + \frac{\cos t(E_a + E_b + m_i) - 1}{(E_a + E_b + m_i)^2} \right], \end{aligned} \quad (22)$$

where $E_{a,b} = (m_{a,b}^2 + \vec{k}_1^2)^{1/2}$.

The crucial difference with respect to the previous model is that, now, Ξ_3 is ultraviolet divergent for any finite value of $t > 0$, as a direct power counting shows. The same statement holds, in principle, for Ξ_4 . To the best of our knowledge, and in spite of the fact that the present model is renormalizable, no cure seems to exist for such divergences. It is not hard to see that the source of the difficulties is related to the fact that the self-energy diagram for the boson is quadratically ultraviolet divergent. Therefore, it seems that if the relevant self-energy diagrams are, at most, logarithmically divergent, then one can obtain a finite survival probability for any $t \geq 0$. Contrarily, if higher order divergences are present, the survival probability does not exist, in the present context of strictly RQFT, for any finite value of $t > 0$, and, then, a different regularization procedure, such as the one introduced in [5]

—which, in a strict sense, lies outside the province of RQFT—should be employed in order to define the physical survival probability for finite times. In any case, one should note that it is formally possible to let $t \rightarrow \infty$ in (22), and employ $[1 - \cos(tq)]/q^2 \rightarrow \pi t \delta(q)$, as in the previous case. This formal limiting process in the divergent (real) quantity Ξ_3 leads to a well defined limit for the survival probability $P(t)$, namely, the actual analogue of Eq. (17) holds as well in the present model: the actual Γ coincides with the total decay rate of the unstable boson in its rest frame.

Going back to the question of quadratic divergences and the non-existence then—in a strict RQFT context—of the survival probability at finite times, it is worth recalling that, in QED, the photon self-energy (vacuum polarization) diagram is quadratically divergent by naive power counting but, actually, it turns out to be only logarithmically divergent (before renormalization) because of gauge invariance (see [8], for instance) —a feature which bears some relationship to the behaviour of the $W^\pm$ and Z^0 self-energies in the Weinberg-Salam model. Therefore one could speculate on whether gauge invariance should play a crucial role to properly analyze time evolution, and in particular the survival probability at finite times of unstable particles in a RQFT context.

As a final comment, we would like to emphasize that our analysis has relied on computations carried out in renormalized perturbation theory. On the other hand, several renormalizable models on RQFT (like the $\lambda\phi^4$ one) are accepted to have a trivial continuum limit [10] in a non-perturbative (renormalization group) framework. In the latter case, the peculiarities of survival amplitudes of unstable particles at finite times, as discussed in this work, would become irrelevant. Here again, (non-abelian) gauge invariance might come to the rescue, at least for QCD: it does make the renormalized theory non trivial in the continuum limit and one could speculate that, upon forcing all ultraviolet divergences to be logarithmic, time evolution would be well defined.

One of us (RFAE) acknowledges the financial support of CICYT (Proyecto AEN97-1693), Spain, and Human Capital and Mobility Programme, European Commission (Contract ERB CHRXCT 940423) Brussels; the other wishes to thank CICYT (Pr. PB94-0194) for the same reason. We are indebted to A.

Dobado, V. Martín Mayol, C. Pérez Martín, F. Ruiz Ruiz and F.J. Ynduráin for discussions and useful information.

References

- [1] L. Fonda, G.C. Ghirardi, A. Rimini, Rep. Prog. Phys. 41 (1978) 587.
- [2] W.H. Itano, D.J. Heinzen, J.J. Bollinger, D.J. Winneland, Phys. Rev. A 41 (1990) 2295.
- [3] P. Facchi, S. Pascazio, Phys. Lett. A 241 (1998) 139.
- [4] J. Glimm, A. Jaffe, in Statistical Mechanics and Quantum Field Theory, edited by C. DeWitt and R. Stora (Gordon and Breach, New York, 1971).
- [5] C. Bernardini, L. Maiani, M. Testa, Phys. Rev. Lett. 71 (1993) 2687;
L. Maiani, M. Testa, Ann. Phys. (NY) 263 (1998) 353.
- [6] N.N. Bogoliubov, D.V. Shirkov, Introduction to the theory of quantized fields (Interscience, New York, 1959), Chap. VI.
- [7] R.F. Alvarez-Estrada, J.L. Sánchez-Gómez, Rev. Mex. Fis. 42 (1996) 216.
- [8] C. Itzykson, J-B. Zuber, Quantum Field Theory (McGraw-Hill, New York, 1980), Chap. 7.
- [9] D. Lurie, Particles and Fields (John Wiley and Sons, New York, 1968), Chap. 6.
- [10] I. Montvay, G. Münster, Quantum Fields on a Lattice (Cambridge University Press, Cambridge, 1994).